arXiv:1411.3248v1 [math.DS] 12 Nov 2014

Bounded solutions of the finite and infinite-dimensional dynamical systems.

Pokutnyi O.A.

Institute of mathematics of NAS of Ukraine, Kiev, Tereshenkivska 3,
E-mail: lenasas@gmail.com

Abstract

Invariant torus are constructed under assumption that the homogeneous system admits an exponential dichotomy on the semi-axes. The main result is closely related with the well-known Palmer's lemma and results of Boichuk A.A., Samoilenko A.M.

Key words: exponential dichotomy, bounded solutions, invariant manifold.

Statement of the problem

Consider the linear inhomogeneous system

$$\frac{d\phi}{dt} = a(\phi), \qquad \frac{dx}{dt} = P(\phi)x + f(\phi), \tag{1}$$

which defined on the direct product of m-dimensional torus $\mathcal{T}_m$ or infinite dimensional torus $\mathcal{T}_\infty$ and the space R^n under assumption that $a(\phi) \in C^1(\mathcal{T}_m)$; $P(\phi)$, $f(\phi) \in C(\mathcal{T}_m)$; $\phi = (\phi_1, \ldots, \phi_m) \in \mathcal{T}_m$; $x = \mathrm{col}(x_1, \ldots, x_n) \in R^n$. It is known, that the problem of exsisting and constructing of invariant torus $x = u(\phi) \in C(\mathcal{T}_m)$, $\phi \in \mathcal{T}_m$, of the system (1) for all $f(\phi) \in C(\mathcal{T}_m)$ can be solved with using Samoilenko-Green function [1, 2]. For uniqueness it is necessary and sufficient for all $f(\phi) \in C(\mathcal{T}_m)$ that homogeneous system has no degenerate torus

$$\frac{d\phi}{dt} = a(\phi), \qquad \frac{dx}{dt} = P(\phi)x. \tag{2}$$

It means, that for all $\phi \in \mathcal{T}_m$ the system

$$\frac{dx}{dt} = P\big(\phi_t(\phi)\big)x \tag{3}$$

is exponentially-dichotomous (e-dichotomous) on the whole axis $R = (-\infty, +\infty)$, i.e. that there is exists projector $C(\phi) = C^2(\phi)$ and not dependent from ϕ, τ constants $K \geq 1$, $\alpha > 0$ such that

$$\begin{gathered} \left\|\Omega_0^t(\phi)C(\phi)\Omega_\tau^0(\phi)\right\| \leq Ke^{-\alpha(t-\tau)}, \quad t \geq \tau, \\ \left\|\Omega_0^t(\phi)(I - C(\phi))\Omega_\tau^0(\phi)\right\| \leq Ke^{-\alpha(\tau-t)}, \quad \tau \geq t, \end{gathered} \tag{4}$$

for all $t, \tau \in R$; $\Omega_\tau^t(\phi)$ $\left(\Omega_\tau^\tau(\phi) = I_n\right)$ — is $(n \times n)$-dimensional fundamental matrix of the system (3); $\phi_t(\phi)-$ is a solution of the Koschi problem $\dot{\phi} = a(\phi)$, $\phi_0(\phi) = \phi$.

Consider the case when the system (3) doesn't have e-dichotomous on the semi-axes R, but e-dichotomous on the semi-axes R_+ and R_- with projectors $C_+(\phi)$ and $C_-(\phi)$ ($C_\pm^2(\phi) = C_\pm(\phi)$) respectively. It means that [3] for the system (3) the next inequalities are true

$$\begin{array}{c} \left\|\Omega_0^t(\phi)C_+(\phi)\Omega_\tau^0(\phi)\right\| \le K_1 e^{-\alpha_1(t-\tau)}, \quad t \ge \tau \quad t,\tau \in \mathbb{R}_+; \\ \left\|\Omega_0^t(\phi)(I - C_+(\phi))\Omega_\tau^0(\phi)\right\| \le K_1 e^{-\alpha_1(\tau-t)}, \quad \tau \ge t, \quad t,\tau \in \mathbb{R}_+; \end{array} \tag{5}$$

$$\begin{array}{c} \left\|\Omega_0^t(\phi)C_-(\phi)\Omega_\tau^0(\phi)\right\| \le K_2 e^{-\alpha_2(t-\tau)}, \quad t \ge \tau, \quad t,\tau \in \mathbb{R}_-; \\ \left\|\Omega_0^t(\phi)(I - C_-(\phi))\Omega_\tau^0(\phi)\right\| \le K_2 e^{-\alpha_2(\tau-t)}, \quad \tau \ge t, \quad t,\tau \in \mathbb{R}_-; \end{array} \tag{6}$$

is the so called critical case. In this article it is necessary and sufficient conditions for the existence of invariant torus $x = u(\phi) \in C(\mathcal{T}_m)$, $\phi \in \mathcal{T}_m$, of the system (1) are obtained in that case. It is necessary and sufficient conditions for inhomogeneity $f(\phi) \in C(\mathcal{T}_m)$, which define invariant manifold are obtained.

Bounded solutions on the whole axis

For fixed $\phi \in \mathcal{T}_m$ general solutions of the problem

$$\frac{dx}{dt} = P\big(\phi_t(\phi)\big)x + f\big(\phi_t(\phi)\big), \tag{7}$$

bounded on the entire semi-axes R_+ и R_-, have the next form

$$x(t,\phi,\xi_1) = \begin{cases} \Omega_0^t(\phi)C_+(\phi)\xi_1 + \int\limits_0^t \Omega_\tau^t(\phi)C_+(\phi_\tau(\phi))f(\phi_\tau(\phi))d\tau - \\ \qquad - \int\limits_t^\infty \Omega_\tau^t(\phi)(I - C_+(\phi_\tau(\phi))f(\phi_\tau(\phi))d\tau, \quad t \ge 0, \\ \Omega_0^t(\phi)(I - C_-(\phi))\xi_1 + \int\limits_{-\infty}^t \Omega_\tau^t(\phi)C_-(\phi_\tau(\phi))f(\phi_\tau(\phi))d\tau - \\ - \int\limits_t^0 \Omega_\tau^t(\phi)\big(I - C_-(\phi_\tau(\phi))f(\phi_\tau(\phi))d\tau, \quad t \le 0, \end{cases} \tag{8}$$

for all bounded f and

$$x(t,\phi,\xi_2)=\begin{cases}\Omega_0^t(\phi)C_+(\phi)\xi_2+\int\limits_0^t\Omega_\tau^t(\phi)(I-C_+(\phi_\tau(\phi))f(\phi_\tau(\phi))d\tau-\\ \qquad-\int\limits_t^\infty\Omega_\tau^t(\phi)(C_+(\phi_\tau(\phi))f(\phi_\tau(\phi))d\tau,\quad t\geq 0,\\ \Omega_0^t(\phi)(I-C_-(\phi))\xi_2+\int\limits_{-\infty}^t\Omega_\tau^t(\phi)(I-C_-(\phi_\tau(\phi))f(\phi_\tau(\phi))d\tau-\\ -\int\limits_t^0\Omega_\tau^t(\phi)\big(C_-(\phi_\tau(\phi))f(\phi_\tau(\phi))d\tau,\quad t\leq 0,\end{cases}\tag{9}$$

but not for all bounded f, where

$$C_+(\phi_\tau(\phi))=\Omega_0^\tau(\phi)C_+(\phi)\Omega_\tau^0(\phi),\qquad C_-(\phi_\tau(\phi))=\Omega_0^\tau(\phi)C_-(\phi)\Omega_\tau^0(\phi).\tag{10}$$

We say about conditions on f below. Here are some well-known relations [1]

$$\begin{array}{ll}\Omega_\tau^t\big(\phi_s(\phi)\big)=\Omega_{\tau+s}^{t+s}(\phi), & \Omega_\tau^t(\phi)\Omega_s^\tau(\phi)=\Omega_s^t(\phi),\\ \big(\Omega_\tau^t(\phi)\big)^{-1}=\Omega_t^\tau(\phi), & \phi_\tau(\phi_s(\phi))=\phi_{\tau+s}(\phi),\end{array}\tag{11}$$

which valid for all $t,\tau,s\in R$, $\phi\in\mathcal{T}_m$.

Solutions (8) and (9) will be bounded on the entire axis R, if the constant vectors $\xi_1=\xi_1(\phi)\in R^n$ and $\xi_2=\xi_2(\phi)\in R^n$ satisfy the next algebraic systems, obtained from the (8) and (9) for $t=0$:

$$\big[C_+(\phi)-(I-C_-(\phi))\big]\xi_1=\int\limits_{-\infty}^0 C_-(\phi)\Omega_\tau^0(\phi)f(\phi_\tau(\phi))d\tau+$$
$$+\int\limits_0^\infty(I-C_+(\phi))\Omega_\tau^0(\phi)f(\phi_\tau(\phi))d\tau.\tag{12}$$

$$\big[C_+(\phi)-(I-C_-(\phi))\big]\xi_2=\int\limits_{-\infty}^0(I-C_-(\phi))\Omega_\tau^0(\phi)f(\phi_\tau(\phi))d\tau+$$
$$+\int\limits_0^\infty(C_+(\phi))\Omega_\tau^0(\phi)f(\phi_\tau(\phi))d\tau.\tag{13}$$

Denote by $D(\phi)=C_+(\phi)-\big(I-C_-(\phi)\big)$ is $(n\times n)$-dimensional matrix, and by $D^+(\phi)$ its Moore-Penrose pseudoinvertible [5]; $P_{N(D)}(\phi)$ and $P_{N(D^*)}(\phi)$ are $(n\times n)$-dimensional orthoprojectors:

$$P^2_{N(D)}(\phi) = P_{N(D)}(\phi) = P^*_{N(D)}(\phi),$$
$$P^2_{N(D^*)}(\phi) = P_{N(D^*)}(\phi) = P^*_{N(D^*)}(\phi),$$

which project R^n onto kernel $N(D) = \ker D(\phi)$ and cokernel $N(D^*) = \ker D^*(\phi)$ of matrix $D(\phi)$;

$$P_{N(D^*)}(\phi) = I - D(\phi)D^+(\phi), \qquad P_{N(D)}(\phi) = I - D^+(\phi)D(\phi).$$

System (12) is solvable if and only if the right part of the system (12) belongs to the orthogonal complement of $N^\perp(D^*(\phi)) = \mathrm{Im}\,(D(\phi))$ of the subspace $N(D^*(\phi))$. It means that

$$P_{N(D^*)}(\phi)\Bigg\{\int\limits_{-\infty}^{0} C_-(\phi)\Omega^0_\tau(\phi)f\big(\phi_\tau(\phi)\big)d\tau +$$
$$+\int\limits_{0}^{\infty}(I - C_+(\phi))\Omega^0_\tau(\phi)f\big(\phi_\tau(\phi)\big)d\tau\Bigg\} = 0. \tag{14}$$

In this case the general solutions of the system (12), bounded on the entire axis R, have the form (8) with constant $\xi_1 = \xi_1(\phi) \in R^n$, which defines from the equation (12) by the rule:

$$\xi_1 = D^+(\phi)\Bigg\{\int\limits_{-\infty}^{0} C_-(\phi)\Omega^0_\tau(\phi)f(\phi_\tau(\phi))d\tau +$$
$$+\int\limits_{0}^{\infty}\big(I - C_+(\phi)\big)\Omega^0_\tau(\phi)f\big(\phi_\tau(\phi)\big)d\tau\Bigg\} + P_{N(D)}(\phi)c, \quad c = c(\phi) \in R^n. \tag{15}$$

Substitute (15) in (8), we obtain that for fixed $\phi \in \mathcal{T}_m$ and inhomogeneity $f(\phi_t(\phi)) \in C(\mathcal{T}_m)$, which satisfies condition (14), solutions, bounded on R of the system (7) have the form

$$x(t,\phi,c) =$$

$$= \Omega_0^t(\phi) \begin{cases} C_+(\phi)P_{N(D)}(\phi)c + \int\limits_0^t C_+(\phi)\Omega_\tau^0(\phi)f(\phi_\tau(\phi))d\tau - \\ \quad - \int\limits_t^\infty \big(I - C_+(\phi)\big)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau + \\ \quad + C_+(\phi)D^+(\phi)\Bigg\{ \int\limits_{-\infty}^0 C_-(\phi)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau + \\ \quad + \int\limits_0^\infty \big(I - C_+(\phi)\big)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau \Bigg\}, \quad t \geq 0, \\ \big(I - C_-(\phi)\big)P_{N(D)}(\phi)c + \int\limits_{-\infty}^t C_-(\phi)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau - \\ \quad - \int\limits_t^0 \big(I - C_-(\phi)\big)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau + \\ \quad + \big(I - C_-(\phi)\big)D^+(\phi)\Bigg\{ \int\limits_{-\infty}^0 C_-(\phi)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau + \\ \quad + \int\limits_0^\infty \big(I - C_+(\phi)\big)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau \Bigg\}, \quad t \leq 0. \end{cases} \tag{16}$$

Since $P_{N(D^*)}(\phi)D(\phi) = P_{N(D^*)}(\phi)\big[C_+(\phi) - (I - C_-(\phi))\big] = 0$, then

$$P_{N(D^*)}(\phi)C_+(\phi) = P_{N(D^*)}(\phi)\big(I - C_-(\phi)\big),$$

and condition (14) is equivalent one of the conditions

$$\begin{aligned} P_{N(D^*)}(\phi) \int\limits_{-\infty}^{+\infty} C_-(\phi)\Omega_\tau^0(\phi)f(\phi_\tau(\phi))d\tau = 0, \\ P_{N(D^*)}(\phi) \int\limits_{-\infty}^{\infty} (I - C_+(\phi))\Omega_\tau^0(\phi)f(\phi_\tau(\phi))d\tau = 0. \end{aligned} \tag{17}$$

Since

$$\Big[C_+(\phi) - \big(I - C_-(\phi)\big)\Big]D^+(\phi) = I - P_{N(D^*)}(\phi),$$

we obtain

$$C_+(\phi)D^+(\phi)\{\ldots\} - I\{\ldots\} = (I - C_-(\phi))D^+(\phi)\{\ldots\},$$

from the condition (14), $\{\ldots\}$ is the expression in (14).

Since $D(\phi)P_{N(D)}(\phi) = \big[C_+(\phi) - (I - C_-(\phi))\big]P_{N(D)}(\phi) = 0$, then

$$C_+(\phi)P_{N(D)}(\phi) = \big(I - C_-(\phi)\big)P_{N(D)}(\phi).$$

Similarly the system (13) is solvable if and only if the right part of system (13) belongs to the orthogonal complement of $N^{\perp}(D^*(\phi)) = \mathrm{Im}\,(D(\phi))$ of the subspace $N(D^*(\phi))$. It means that

$$P_{N(D^*)}(\phi)\Bigg\{\int\limits_{-\infty}^{0}(I - C_-(\phi))\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau + \int\limits_{0}^{\infty}(C_+(\phi))\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau\Bigg\} = 0. \tag{18}$$

In this case the general solutions of the system (13), bounded on the entire axis R, have the form (9) with constant $\xi_2 = \xi_2(\phi) \in R^n$, which defines from the equation (13) by the rule:

$$\xi_2 = D^+(\phi)\Bigg\{\int\limits_{-\infty}^{0}(I - C_-(\phi))\Omega_\tau^0(\phi)f(\phi_\tau(\phi))d\tau + \int\limits_{0}^{\infty}\big(C_+(\phi)\big)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau\Bigg\} + P_{N(D)}(\phi)c, \quad c = c(\phi) \in R^n. \tag{19}$$

Substitute (19) in (9), we obtain that for fixed $\phi \in \mathcal{T}_m$ and inhomogeneity $f(\phi_t(\phi)) \in C(\mathcal{T}_m)$, which satisfies condition (18), solutions, bounded on R of the system (7) have the form

$$x(t,\phi,c) =$$

$$= \Omega_0^t(\phi) \begin{cases} C_+(\phi)P_{N(D)}(\phi)c + \int\limits_0^t (I - C_+(\phi))\Omega_\tau^0(\phi)f(\phi_\tau(\phi))d\tau - \\ \quad - \int\limits_t^\infty \big(C_+(\phi)\big)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau + \\ \quad + C_+(\phi)D^+(\phi)\Bigg\{ \int\limits_{-\infty}^0 (I - C_-(\phi))\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau + \\ \quad + \int\limits_0^\infty \big(C_+(\phi)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau \Bigg\}, \quad t \geq 0, \\ \big(I - C_-(\phi)\big)P_{N(D)}(\phi)c + \int\limits_{-\infty}^t (I - C_-(\phi))\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau - \\ \quad - \int\limits_t^0 \big(C_-(\phi)\big)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau + \\ \quad + \big(I - C_-(\phi)\big)D^+(\phi)\Bigg\{ \int\limits_{-\infty}^0 (I - C_-(\phi))\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau + \\ \quad + \int\limits_0^\infty \big(C_+(\phi)\big)\Big)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau \Bigg\}, \quad t \leq 0. \end{cases} \tag{20}$$

Since $P_{N(D^*)}(\phi)D(\phi) = P_{N(D^*)}(\phi)\big[I - C_+(\phi) + C_-(\phi))\big] = 0$, then

$$P_{N(D^*)}(\phi)(I - C_+(\phi)) = P_{N(D^*)}(\phi)\big(C_-(\phi)\big),$$

and condition (18) is equivalent one of the conditions

$$\begin{aligned} P_{N(D^*)}(\phi) \int\limits_{-\infty}^{+\infty} (I - C_-(\phi))\Omega_\tau^0(\phi)f(\phi_\tau(\phi))d\tau = 0, \\ P_{N(D^*)}(\phi) \int\limits_{-\infty}^{\infty} (C_+(\phi))\Omega_\tau^0(\phi)f(\phi_\tau(\phi))d\tau = 0. \end{aligned} \tag{21}$$

Since

$$\Big[I - C_+(\phi) + C_-(\phi))\Big]D^+(\phi) = I - P_{N(D^*)}(\phi),$$

we obtain

$$(I - C_+(\phi))D^+(\phi)\{\ldots\} - I\{\ldots\} = (C_-(\phi))D^+(\phi)\{\ldots\},$$

from the condition (18), $\{\ldots\}$ is expression in (18).

Since $D(\phi)P_{N(D)}(\phi) = \big[C_+(\phi) - (I - C_-(\phi))\big]P_{N(D)}(\phi) = 0$, then

$$(I - C_+(\phi))P_{N(D)}(\phi) = \big(C_-(\phi)\big)P_{N(D)}(\phi).$$

Consider the case, when homogeneous system (3) does not have bounded and unbounded solutions

$$C_+(\phi)P_{N(D)}(\phi) = \big(I - C_-(\phi)\big)P_{N(D)}(\phi) = 0.$$

Then (16) and (20) we can rewrite in the form

$$x(t,\phi) = (G_t(f))(\phi), \tag{22}$$

$$\big(G_t(f)\big)(\phi) = \Omega_0^t(\phi)\begin{cases} \int\limits_0^t C_+(\phi)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau - \\ \quad - \int\limits_t^\infty \big(I - C_+(\phi)\big)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau + \\ \quad + C_+(\phi)D^+(\phi)\Big\{ \int\limits_{-\infty}^0 C_-(\phi)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau + \\ \quad + \int\limits_0^\infty \big(I - C_+(\phi)\big)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau \Big\}, \quad t \geq 0, \\ \int\limits_{-\infty}^t C_-(\phi)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau - \\ \quad - \int\limits_t^0 \big(I - C_-(\phi)\big)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau + \\ \quad + \big[C_+(\phi)D^+(\phi) - I\big]\Big\{ \int\limits_{-\infty}^0 C_-(\phi)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau + \\ \quad + \int\limits_0^\infty \big(I - C_+(\phi)\big)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau \Big\}, \quad t \leq 0, \end{cases}$$

and in the second case

$$\big(G_t(f)\big)(\phi)=\Omega_0^t(\phi)\begin{cases}\int\limits_0^t(I-C_+(\phi))\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau-\\ \qquad-\int\limits_t^\infty\big(C_+(\phi)\big)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau+\\ \qquad+C_+(\phi)D^+(\phi)\Big\{\int\limits_{-\infty}^0(I-C_-(\phi))\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau\\ \qquad+\int\limits_0^\infty\big(C_+(\phi)\big)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau\Big\},\quad t\geq 0,\\ \int\limits_{-\infty}^t(I-C_-(\phi))\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau-\\ \qquad-\int\limits_t^0\big(C_-(\phi)\big)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau+\\ \qquad+\big[(I-C_+(\phi))D^+(\phi)-I\big]\\ \Big\{\int\limits_{-\infty}^0(I-C_-(\phi))\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau+\\ \qquad+\int\limits_0^\infty\big(C_+(\phi)\big)\Omega_\tau^0(\phi)f\big(\phi_\tau(\phi)\big)d\tau\Big\},\quad t\leq 0,\end{cases}$$

will be called generalized Green's operator of the problem about invariant torus of the system (1).

Under conditions (15), (19) solutions, bounded on R, of the system (7) for fixed $\phi\in\mathcal{T}_m$ have the form (22).

We show, that the expression

$$x(0,\phi)=u(\phi)=\big(G_0(f)\big)(\phi),\tag{23}$$

which obtained from (22) for $t=0$, define for all $\phi\in\mathcal{T}_m$ invariant torus of the system (1).

Criterion of existence of invariant torus of nonhomogeneous system

As shown below, under conditions

$$P_{N(D^*)}(\phi)\int\limits_{-\infty}^{+\infty}C_-(\phi)\Omega_\tau^0(\phi)f(\phi_\tau(\phi))d\tau=0,\tag{24}$$

$$P_{N(D^*)}(\phi)\int\limits_{-\infty}^{+\infty}(I-C_-(\phi))\Omega_\tau^0(\phi)f(\phi_\tau(\phi))d\tau=0,\tag{25}$$

the nonhomogeneous system (7) have bounded solutions on R in the form (22) for fixed $\phi \in \mathcal{T}_m$. Conditions (24) and (25) on solutions $\phi_t(\phi)$ define invariant set. Substitute $\phi_t(\phi)$ instead of ϕ and show that conditions (24) and (25) hold for all $t \in R$ and $\phi \in \mathcal{T}_m$. From the relations for $D(\phi) = C_+(\phi) - \big(I - C_-(\phi)\big)$ we obtain the next equality

$$D\big(\phi_t(\phi)\big) = \Omega_0^t(\phi) D(\phi) \Omega_t^0(\phi) \quad \forall t \in R, \quad \forall \phi \in \mathcal{T}_m. \tag{26}$$

Direct check shows, that for all $t \in R$ and $\phi \in \mathcal{T}_m$ the matrix

$$D^-\big(\phi_t(\phi)\big) = \big[\Omega_0^t(\phi) D(\phi) \Omega_t^0(\phi)\big]^- = \Omega_0^t(\phi) D^-(\phi) \Omega_t^0(\phi) \tag{27}$$

is generalized-invertible to the matrix $D\big(\phi_t(\phi)\big)$ and satisfies the next relations [5]

$$\begin{aligned} D^-\big(\phi_t(\phi)\big) D\big(\phi_t(\phi)\big) D^-\big(\phi_t(\phi)\big) &= D^-\big(\phi_t(\phi)\big), \\ D\big(\phi_t(\phi)\big) D^-\big(\phi_t(\phi)\big) D\big(\phi_t(\phi)\big) &= D\big(\phi_t(\phi)\big). \end{aligned} \tag{28}$$

From the conditions

$$\begin{aligned} D\big(\phi_t(\phi)\big) D^-\big(\phi_t(\phi)\big) &= I - P_{N(D)}\big(\phi_t(\phi)\big), \\ D^-\big(\phi_t(\phi)\big) D\big(\phi_t(\phi)\big) &= I - P_{N(D^*)}\big(\phi_t(\phi)\big) \end{aligned}$$

we get expressions for projectors $P_{N(D)}(\phi_t(\phi))$ and $P_{N(D^*)}(\phi_t(\phi))$ onto kernel and cokernel of matrix $D(\phi)$ on solutions $\phi_t(\phi)$ of the respectively Koschi problem for all $t \in R$ and $\phi \in \mathcal{T}_m$:

$$\begin{aligned} P_{N(D)}\big(\phi_t(\phi)\big) &= \Omega_0^t(\phi) P_{N(D)}(\phi) \Omega_t^0(\phi) = \Omega_0^t(\phi) \big[I - D^-(\phi) D(\phi)\big] \Omega_t^0(\phi), \\ P_{N(D^*)}\big(\phi_t(\phi)\big) &= \Omega_0^t(\phi) P_{N(D^*)}(\phi) \Omega_t^0(\phi) = \Omega_0^t(\phi) \big[I - D(\phi) D^-(\phi)\big] \Omega_t^0(\phi). \end{aligned} \tag{29}$$

We can choose that $D^-(\phi) = D^+(\phi)$. In that case projectors $P_{N(D)}(\phi)$ and $P_{N(D^*)}(\phi)$ will be orthoprojectors.

For all $t \in R$ and $\phi \in \mathcal{T}_m$ we have

$$P_{N(D^*)}\big(\phi_t(\phi)\big) \int\limits_{-\infty}^{+\infty} C_-\big(\phi_t(\phi)\big) \Omega_\tau^0\big(\phi_t(\phi)\big) f\big(\phi_\tau\big(\phi_t(\phi)\big)\big) d\tau =$$

$$= \Omega_0^t(\phi) P_{N(D^*)}(\phi) \int\limits_{-\infty}^{+\infty} C_-(\phi) \Omega_{\tau+t}^0(\phi) f(\phi_{\tau+t}(\phi)) d\tau = 0,$$

and

$$\Omega_0^t(\phi) P_{N(D^*)}(\phi) \int\limits_{-\infty}^{+\infty} (I - C_-(\phi)) \Omega_{\tau+t}^0(\phi) f(\phi_{\tau+t}(\phi)) d\tau = 0,$$

From the conditions (10), (12), (29) we have

$$u\big(\phi_t(\phi)\big) = (G_0(f))\big(\phi_t(\phi)\big) = \big(G_t(f)\big)(\phi)$$

for all $t \in R$ and $\phi \in \mathcal{T}_m$. It shows that $u\big(\phi_t(\phi)\big) \in C^1(\mathcal{T}_m)$, and the set $u(\phi)$ defines invariant torus of the system (1).

In a such way we have the following theorem.

Theorem. *Let the system* (3) *is e-dichotomous on both semi-axes* R_+ *u* R_- *with projectors* $C_\pm(\phi)$, *which satisfy the next equalities for* $\phi_t(\phi)$

$$C_\pm\big(\phi_t(\phi)\big) = \Omega_0^t(\phi)C_\pm(\phi)\Omega_t^0(\phi), \quad C_\pm^2(\phi) = C_\pm(\phi).$$

The system (1) *has invariant torus if and only if nonhomogeneity* $f(\phi) \in C(\mathcal{T}_m)$ *satisfies conditions* (24), (25). *If the homogeneous system* (3) *does not have bounded and unbounded solutions, i.e. the next condition is true*

$$C_+(\phi)P_{N(D)}(\phi) = \big(I - C_-(\phi)\big)P_{N(D)}(\phi) = 0,$$

then expression

$$u(\phi) = \big(G_0(f)\big)(\phi)$$

which obtain from (22), *for* $t = 0$, *defines for all* $\phi \in \mathcal{T}_m$ *invariant torus of system* (1).

Examples. Consider the problem of the existence of invariant manifold of the system

$$\dot{\varphi} = 1, \tag{30}$$

$$\tag{31}$$

$$\dot{x}_1(t) = \mathrm{th}\,(\varphi)x(t) + f_1(\varphi). \tag{32}$$

$$\dot{x}_2(t) = -\mathrm{th}\,(\varphi)x_2(t) + f_2(\varphi). \tag{33}$$

This system has the following characteristics:

$$\Omega_0^t(\varphi) = \begin{pmatrix} \frac{ch(\varphi_t(\phi))}{ch\varphi} & 0 \\ 0 & \frac{ch\varphi}{ch\varphi_t(\varphi)} \end{pmatrix},$$

$$C_+(\varphi) = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, C_-(\varphi) = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix},$$

$$D(\varphi) = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} = D^+(\varphi),$$

$$P_{N(D)}(\varphi) = P_{N(D^*)}(\varphi) = I.$$

In this case the first condition of solvability have the form

$$\int_{-\infty}^{+\infty} \frac{ch\varphi}{ch(\varphi_\tau(\varphi))} f_1(\varphi_\tau(\varphi))d\tau = 0,$$

and under this condition the system has the invariant torus in the form

$$u_1(\varphi) = \begin{pmatrix} -\int_0^{+\infty} \frac{ch(\varphi) f_1(\varphi_\tau(\varphi))}{ch(\varphi_\tau(\varphi))} d\tau \\ 0 \end{pmatrix},$$

and under the second condition of solvability

$$\int_{-\infty}^{+\infty} ch(\varphi_\tau(\varphi)) f_2(\varphi_\tau(\varphi)) d\tau = 0$$

the system has the invariant torus

$$u_2(\varphi) = \begin{pmatrix} 0 \\ -\int_0^{+\infty} \frac{ch(\varphi_\tau(\varphi)) f_2(\varphi_\tau(\varphi))}{ch(\varphi)} d\tau \end{pmatrix}.$$

If, for example $f_1(\varphi) = \frac{sh(\varphi)}{ch^3(\varphi)}$ and $f_2(\varphi) = \frac{sh(\varphi)}{ch^4(\varphi)}$, then the given system has invariant torus in the glued form

$$u(\varphi) = \begin{pmatrix} -\frac{1}{3ch^2(\varphi)} \\ -\frac{1}{2ch^3(\varphi)} \end{pmatrix}.$$

We mention that given theory works in the case of infinite dimensional space. Here is an example. Consider the countable system of differential equations in the space $BC(\mathbb{R}, l_2)$ or $BC(\mathbb{R}, l_{2loc})$ in the next form

$$\dot{\varphi}(t) = 1,$$

$$\frac{dx(t)}{dt} = P(\varphi_t(\varphi))x(t) + f(\varphi_t(\varphi)),$$

where

$$x(t) = (x_1(t), x_2(t), ...) \in l_2 \quad \text{for all} \quad t,$$

and

$$f(\varphi) = (f_1(\varphi), f_2(\varphi), ...),$$

where

$$P(\varphi) = diag\{th(\varphi), th(\varphi), -th(\varphi), -th(\varphi), -th(\varphi), ...\}.$$

$$x(t) = (x_1(t), x_2(t), ...), f(t) = (f_1(t), f_2(t), ...) \in BC(\mathbb{R}, l_2).$$

Here $BC(\mathbb{R}, l_2)$ and $BC(\mathbb{R}, l_{2loc})$ are the spaces of bounded and continuous on the whole axis functions with values in l_2 or l_{2loc}.

Matriciant of the system has the form

$$\Omega_0^t(\varphi) = diag\{\frac{ch(\varphi_t(\varphi))}{ch\varphi}, \frac{ch(\varphi_t(\varphi))}{ch\varphi}, \frac{ch\varphi}{ch(\varphi_t(\varphi))}, \frac{ch\varphi}{ch(\varphi_t(\varphi))}, \frac{ch\varphi}{ch(\varphi_t(\varphi))}, ...\}.$$

Projectors have the form

$$C_+(\varphi) = diag\{0, 0, 1, 1, ...\}, C_-(\varphi) = diag\{1, 1, 0, 0, ...\}.$$

Matrixes $D(\varphi) = D^+(\varphi) = 0$, and $P_{N(D)} = P_{N(D^*)} = I$, where I is the identity matrix. Condition of the solvability for the first type of torus has the form

$$\int_{-\infty}^{+\infty} \frac{f_i(\varphi_\tau(\varphi))}{ch(\varphi_\tau(\varphi))} d\tau = 0,\ i = 1, 2,$$

and invariant torus has the form

$$x = u_1(\varphi) = (-\int_0^{+\infty} \frac{ch(\varphi) f_1(\varphi_\tau(\varphi))}{ch(\varphi_\tau(\varphi))} d\tau, -\int_0^{+\infty} \frac{ch(\varphi) f_2(\varphi_\tau(\varphi))}{ch(\varphi_\tau(\varphi))} d\tau, 0, ...).$$

Condition of the solvability for the second type has the form

$$\int_{-\infty}^{+\infty} f_i(\varphi_\tau(\varphi)) ch(\varphi_\tau(\varphi)) d\tau = 0, i \geq 3,$$

and invariant torus has the form

$$x = u_2(\varphi) = (0, 0, -\int_0^{+\infty} \frac{f_3(\varphi_\tau(\varphi)) ch(\varphi_\tau(\varphi))}{ch(\varphi)} d\tau, ..., -\int_0^{+\infty} \frac{f_i(\varphi_\tau(\varphi)) ch(\varphi_\tau(\varphi))}{ch(\varphi)} d\tau, ...),$$

or, in the glued form

$$x = u(\varphi) = (-\int_0^{+\infty} \frac{ch(\varphi) f_1(\varphi_\tau(\varphi))}{ch(\varphi_\tau(\varphi))} d\tau, -\int_0^{+\infty} \frac{ch(\varphi) f_2(\varphi_\tau(\varphi))}{ch(\varphi_\tau(\varphi))} d\tau,$$

$$-\int_0^{+\infty} \frac{f_3(\varphi_\tau(\varphi)) ch(\varphi_\tau(\varphi))}{ch(\varphi)} d\tau, ..., -\int_0^{+\infty} \frac{f_i(\varphi_\tau(\varphi)) ch(\varphi_\tau(\varphi))}{ch(\varphi)} d\tau, ...).$$

If, for example, $f_i(\varphi) = \frac{sh\varphi}{ch^{i+2}(\varphi)}$, $i \geq 1$, then we have

$$x = u_1(\varphi) = (-\frac{1}{2ch^2(\varphi)}, -\frac{1}{3ch^3(\varphi)}, 0, ...),$$

and

$$x = u_2(\varphi) = (0, 0, -\frac{1}{3ch^4(\varphi)}, ..., -\frac{1}{ich^{i+1}(\varphi)}, ...),$$

or in the glued form

$$x = u(\varphi) = (-\frac{1}{2ch^2(\varphi)}, -\frac{1}{3ch^3(\varphi)}, -\frac{1}{3ch^4(\varphi)}, ..., -\frac{1}{ich^{i+1}(\varphi)}, ...).$$

Here is denotions as in [8].

Bibliography

[1] *Samoilenko A.M.* Elements of mathematical theory of multifrequency oscillations. – M.: Science, 1987. – 304 p. (in russian).

[2] *Mitropolsky Yu.O., Samoilenko A.M., Kulik V.L.* Investigation of dichotomy of linear system of differential equations with Lyapunov functions. – Kiev, 1990. – 270 p. (in russian)

[3] *Palmer K. J.* Exponential dichotomies and transversal homoclinic points // J. Different. Equat. – 1984. – **55**. – P. 225 – 256.

[4] *Boichuk A. A.* Solutions of weakly nonlinear differential equations bounded on the whole line // Nonlinear Oscillations. – 1999. – **2**, No 1. – P. 3 – 10.

[5] *Boichuk A. A., Samoilenko A. M.* Generalized inverse operators and fredholm boundary value problems. – Utrecht; Boston: VSP, 2004. – 317 p.

[6] *Boichuk A.A.* Condition of existence of unique Green-Samoilenko function of the invariant torus problem // Ukrainian Math. Journ. – 2001. – **53**, No4. – p. 556 – 559.

[7] *Boichuk A.* Bounded solutions of differential equations in Banach space // Colloq. Different. and Difference Equat. dedicat. Prof. Jaroslav Kurzweil 80-th Birthday: Abstrs (Brno, Czech Republic, Sept. 5 – 8, 2006). – P. 35.

[8] *Boichuk A.A.* Criterion of existence of unique invariant torus of linear extensrion dynamical systems. – Ukrainian Math. Journal, 2007, V.59, №1. – p.3 – 13.

Institute of mathematics of NAS of Ukraine, Kiev, 01601, Tereshenkivska str. 3,
lenasas@gmail.com